

Examples of PDEs all whose points are characteristic

M. Eugenia Rosado
Departamento de Matemática Aplicada, ETSAM, UPM
Madrid, Spain

June 2011

Invariant Lagrangians on the bundle of linear frames

Invariant Lagrangians on the bundle of linear frames

M smooth manifold, $m = \dim M$.

Invariant Lagrangians on the bundle of linear frames

M smooth manifold, $m = \dim M$.

$\pi: FM \rightarrow M$ the bundle of linear frames of M .

Invariant Lagrangians on the bundle of linear frames

M smooth manifold, $m = \dim M$.

$\pi: FM \rightarrow M$ the bundle of linear frames of M .

\mathcal{L} Lagrangian function on $J^1(FM)$.

Invariant Lagrangians on the bundle of linear frames

M smooth manifold, $m = \dim M$.

$\pi: FM \rightarrow M$ the bundle of linear frames of M .

\mathcal{L} Lagrangian function on $J^1(FM)$.

$$\mathcal{L} \text{ is } \text{Diff}(M)\text{-invariant} \iff \mathcal{L} \circ J^1(\tilde{\varphi}) = \mathcal{L}, \quad \forall \varphi \in \text{Diff}(M).$$

Invariant Lagrangians on the bundle of linear frames

M smooth manifold, $m = \dim M$.

$\pi: FM \rightarrow M$ the bundle of linear frames of M .

\mathcal{L} Lagrangian function on $J^1(FM)$.

$$\mathcal{L} \text{ is } \text{Diff}(M)\text{-invariant} \iff \mathcal{L} \circ J^1(\tilde{\varphi}) = \mathcal{L}, \quad \forall \varphi \in \text{Diff}(M).$$

Natural lifts of φ :

$$\begin{aligned}\tilde{\varphi}: FM &\rightarrow FM, \\ J^1(\tilde{\varphi}): J^1(FM) &\rightarrow J^1(FM).\end{aligned}$$

Invariant Lagrangians on the bundle of linear frames

M smooth manifold, $m = \dim M$.

$\pi: FM \rightarrow M$ the bundle of linear frames of M .

\mathcal{L} Lagrangian function on $J^1(FM)$.

$$\mathcal{L} \text{ is } \text{Diff}(M)\text{-invariant} \iff \mathcal{L} \circ J^1(\tilde{\varphi}) = \mathcal{L}, \quad \forall \varphi \in \text{Diff}(M).$$

Natural lifts of φ :

$$\begin{aligned} \tilde{\varphi}: FM &\rightarrow FM, \\ J^1(\tilde{\varphi}): J^1(FM) &\rightarrow J^1(FM). \end{aligned}$$

$$\mathcal{L} \text{ is } \mathfrak{X}(M)\text{-invariant} \iff \tilde{X}^{(1)}(\mathcal{L}) = 0, \quad \forall X \in \mathfrak{X}(M).$$

Invariant Lagrangians on the bundle of linear frames

M smooth manifold, $m = \dim M$.

$\pi: FM \rightarrow M$ the bundle of linear frames of M .

\mathcal{L} Lagrangian function on $J^1(FM)$.

$$\mathcal{L} \text{ is } \text{Diff}(M)\text{-invariant} \iff \mathcal{L} \circ J^1(\tilde{\varphi}) = \mathcal{L}, \quad \forall \varphi \in \text{Diff}(M).$$

Natural lifts of φ :

$$\begin{aligned}\tilde{\varphi}: FM &\rightarrow FM, \\ J^1(\tilde{\varphi}): J^1(FM) &\rightarrow J^1(FM).\end{aligned}$$

$$\mathcal{L} \text{ is } \mathfrak{X}(M)\text{-invariant} \iff \tilde{X}^{(1)}(\mathcal{L}) = 0, \quad \forall X \in \mathfrak{X}(M).$$

\tilde{X} natural lift of a vector field $X \in \mathfrak{X}(M)$ to FM ,

$\tilde{X}^{(1)}$ natural lift of \tilde{X} to $J^1(FM)$ by infinitesimal contact transformations.

Basis of invariant Lagrangians

Basis of invariant Lagrangians

$s = (X_1, \dots, X_m)$ linear frame with dual coframe $(\omega^1, \dots, \omega^m)$.

Basis of invariant Lagrangians

$s = (X_1, \dots, X_m)$ linear frame with dual coframe $(\omega^1, \dots, \omega^m)$. We consider:

$$\begin{aligned}\mathcal{L}_{jk}^i &: J^1(FM) \rightarrow \mathbb{R}, \quad j < k, \\ \mathcal{L}_{jk}^i(j_x^1 s) &= \omega^i([X_j, X_k])(x).\end{aligned}$$

Basis of invariant Lagrangians

$s = (X_1, \dots, X_m)$ linear frame with dual coframe $(\omega^1, \dots, \omega^m)$. We consider:

$$\begin{aligned}\mathcal{L}_{jk}^i &: J^1(FM) \rightarrow \mathbb{R}, \quad j < k, \\ \mathcal{L}_{jk}^i(j_x^1 s) &= \omega^i([X_j, X_k])(x).\end{aligned}$$

The functions \mathcal{L}_{jk}^i are $\text{Diff}M$ -invariant and functionally independent.

Basis of invariant Lagrangians

$s = (X_1, \dots, X_m)$ linear frame with dual coframe $(\omega^1, \dots, \omega^m)$. We consider:

$$\begin{aligned}\mathcal{L}_{jk}^i &: J^1(FM) \rightarrow \mathbb{R}, \quad j < k, \\ \mathcal{L}_{jk}^i(j_x^1 s) &= \omega^i([X_j, X_k])(x).\end{aligned}$$

The functions \mathcal{L}_{jk}^i are $\text{Diff}M$ -invariant and functionally independent.

If $\dim M = m \geq 3$, the Lagrangians \mathcal{L}_{jk}^i define two types of non equivalent variational problems according to $i \in \{j, k\}$ or $i \in \{j, k\}^c$.

Basis of invariant Lagrangians

$s = (X_1, \dots, X_m)$ linear frame with dual coframe $(\omega^1, \dots, \omega^m)$. We consider:

$$\begin{aligned}\mathcal{L}_{jk}^i &: J^1(FM) \rightarrow \mathbb{R}, \quad j < k, \\ \mathcal{L}_{jk}^i(j_x^1 s) &= \omega^i([X_j, X_k])(x).\end{aligned}$$

The functions \mathcal{L}_{jk}^i are $\text{Diff}M$ -invariant and functionally independent.

If $\dim M = m \geq 3$, the Lagrangians \mathcal{L}_{jk}^i define two types of non equivalent variational problems according to $i \in \{j, k\}$ or $i \in \{j, k\}^c$.

We only consider $\mathcal{L}_{12}^1, \mathcal{L}_{23}^1$, $\dim M = m \geq 3$.

Euler-Lagrange equations

Euler-Lagrange equations

A linear frame $s = (X_1, \dots, X_m)$ with dual coframe $(\omega^1, \dots, \omega^m)$ is an extremal of \mathcal{L}_{23}^1 if and only if the following $3(m-2)$ equations hold:

Euler-Lagrange equations

A linear frame $s = (X_1, \dots, X_m)$ with dual coframe $(\omega^1, \dots, \omega^m)$ is an extremal of \mathcal{L}_{23}^1 if and only if the following $3(m-2)$ equations hold:

$$0 = \omega^1([X_3, X_i]),$$

$$0 = \omega^1([X_2, X_i]),$$

$$0 = 2\omega^1([X_1, X_3]) + \sum_{h=4}^m \omega^h([X_h, X_3]),$$

$$0 = 2\omega^1([X_1, X_2]) + \sum_{h=4}^m \omega^h([X_h, X_2]),$$

$$0 = \omega^1([X_2, X_3]),$$

$$0 = \omega^i([X_2, X_3]),$$

Euler-Lagrange equations

A linear frame $s = (X_1, \dots, X_m)$ with dual coframe $(\omega^1, \dots, \omega^m)$ is an extremal of \mathcal{L}_{23}^1 if and only if the following $3(m-2)$ equations hold:

$$0 = \omega^1([X_3, X_i]),$$

$$0 = \omega^1([X_2, X_i]),$$

$$0 = 2\omega^1([X_1, X_3]) + \sum_{h=4}^m \omega^h([X_h, X_3]),$$

$$0 = 2\omega^1([X_1, X_2]) + \sum_{h=4}^m \omega^h([X_h, X_2]),$$

$$0 = \omega^1([X_2, X_3]),$$

$$0 = \omega^i([X_2, X_3]),$$

where $4 \leq i \leq m$.

Euler-Lagrange equations

A linear frame $s = (X_1, \dots, X_m)$ with dual coframe $(\omega^1, \dots, \omega^m)$ is an extremal of \mathcal{L}_{23}^1 if and only if the following $3(m-2)$ equations hold:

$$0 = \omega^1([X_3, X_i]),$$

$$0 = \omega^1([X_2, X_i]),$$

$$0 = 2\omega^1([X_1, X_3]) + \sum_{h=4}^m \omega^h([X_h, X_3]),$$

$$0 = 2\omega^1([X_1, X_2]) + \sum_{h=4}^m \omega^h([X_h, X_2]),$$

$$0 = \omega^1([X_2, X_3]),$$

$$0 = \omega^i([X_2, X_3]),$$

where $4 \leq i \leq m$.

For the Lagrangian \mathcal{L}_{12}^1 we obtain a similar result with $3(m-1)$ equations.

Euler-Lagrange equations: Local expression

Euler-Lagrange equations: Local expression

Coordinate domain $(U; x^i)$ in M . Section: $s = (X_1, \dots, X_m)$ with

$$X_j = f_j^i \frac{\partial}{\partial x^i}, \quad f_j^i \in C^\infty(U).$$

Euler-Lagrange equations: Local expression

Coordinate domain $(U; x^i)$ in M . Section: $s = (X_1, \dots, X_m)$ with

$$X_j = f_j^i \frac{\partial}{\partial x^i}, \quad f_j^i \in C^\infty(U).$$

Euler-Lagrange equations of \mathcal{L}_{23}^1 :

$$\left(f_a^k \frac{\partial f_i^h}{\partial x^k} - f_i^k \frac{\partial f_a^h}{\partial x^k} \right) f_h^{1} = 0, \quad \begin{array}{l} 4 \leq i \leq m, \\ a = 2, 3, \end{array}$$

$$2 \left(f_1^k \frac{\partial f_a^h}{\partial x^k} - f_a^k \frac{\partial f_1^h}{\partial x^k} \right) f_h^{1} + \sum_{b=4}^m \left(f_b^k \frac{\partial f_a^h}{\partial x^k} - f_a^k \frac{\partial f_b^h}{\partial x^k} \right) f_h^b = 0, \quad a = 2, 3,$$

$$\left(f_2^k \frac{\partial f_3^h}{\partial x^k} - f_3^k \frac{\partial f_2^h}{\partial x^k} \right) f_h^j = 0, \quad j \neq 2, 3.$$

PDEs of extremals of \mathcal{L}_{23}^1

- ① The morphism $\varphi: J^1(FM) \rightarrow M \times \mathbb{R}^{3(m-2)}$ of fibred manifolds over M given by:

$$\begin{aligned}\varphi^{i-3}(j_x^1 s) &= \omega^1([X_3, X_i]_x), \\ \varphi^{m-6+i}(j_x^1 s) &= \omega^1([X_2, X_i]_x), \\ \varphi^{2m-5}(j_x^1 s) &= 2\omega^1([X_1, X_3]_x) + \sum_{h=4}^m \omega^h([X_h, X_3]_x), \\ \varphi^{2m-4}(j_x^1 s) &= 2\omega^1([X_1, X_2]_x) + \sum_{h=4}^m \omega^h([X_h, X_2]_x), \\ \varphi^{2m-3}(j_x^1 s) &= \omega^1([X_2, X_3]_x), \\ \varphi^{2m-6+i}(j_x^1 s) &= \omega^i([X_2, X_3]_x),\end{aligned}$$

where $4 \leq i \leq m$,

- ① The morphism $\varphi: J^1(FM) \rightarrow M \times \mathbb{R}^{3(m-2)}$ of fibred manifolds over M given by:

$$\begin{aligned}\varphi^{i-3}(j_x^1 s) &= \omega^1([X_3, X_i]_x), \\ \varphi^{m-6+i}(j_x^1 s) &= \omega^1([X_2, X_i]_x), \\ \varphi^{2m-5}(j_x^1 s) &= 2\omega^1([X_1, X_3]_x) + \sum_{h=4}^m \omega^h([X_h, X_3]_x), \\ \varphi^{2m-4}(j_x^1 s) &= 2\omega^1([X_1, X_2]_x) + \sum_{h=4}^m \omega^h([X_h, X_2]_x), \\ \varphi^{2m-3}(j_x^1 s) &= \omega^1([X_2, X_3]_x), \\ \varphi^{2m-6+i}(j_x^1 s) &= \omega^i([X_2, X_3]_x),\end{aligned}$$

where $4 \leq i \leq m$, is quasilinear.

PDEs of extremals of \mathcal{L}_{23}^1

- ① The morphism $\varphi: J^1(FM) \rightarrow M \times \mathbb{R}^{3(m-2)}$ of fibred manifolds over M given by:

$$\begin{aligned}\varphi^{i-3}(j_x^1 s) &= \omega^1([X_3, X_i]_x), \\ \varphi^{m-6+i}(j_x^1 s) &= \omega^1([X_2, X_i]_x), \\ \varphi^{2m-5}(j_x^1 s) &= 2\omega^1([X_1, X_3]_x) + \sum_{h=4}^m \omega^h([X_h, X_3]_x), \\ \varphi^{2m-4}(j_x^1 s) &= 2\omega^1([X_1, X_2]_x) + \sum_{h=4}^m \omega^h([X_h, X_2]_x), \\ \varphi^{2m-3}(j_x^1 s) &= \omega^1([X_2, X_3]_x), \\ \varphi^{2m-6+i}(j_x^1 s) &= \omega^i([X_2, X_3]_x),\end{aligned}$$

where $4 \leq i \leq m$, is quasilinear.

- ② The equations of the extremals of \mathcal{L}_{23}^1 are given by $R^1 = \varphi^{-1}(M \times \{0\})$.

PDEs of extremals of \mathcal{L}_{23}^1

- ① The morphism $\varphi: J^1(FM) \rightarrow M \times \mathbb{R}^{3(m-2)}$ of fibred manifolds over M given by:

$$\begin{aligned}\varphi^{i-3}(j_x^1 s) &= \omega^1([X_3, X_i]_x), \\ \varphi^{m-6+i}(j_x^1 s) &= \omega^1([X_2, X_i]_x), \\ \varphi^{2m-5}(j_x^1 s) &= 2\omega^1([X_1, X_3]_x) + \sum_{h=4}^m \omega^h([X_h, X_3]_x), \\ \varphi^{2m-4}(j_x^1 s) &= 2\omega^1([X_1, X_2]_x) + \sum_{h=4}^m \omega^h([X_h, X_2]_x), \\ \varphi^{2m-3}(j_x^1 s) &= \omega^1([X_2, X_3]_x), \\ \varphi^{2m-6+i}(j_x^1 s) &= \omega^i([X_2, X_3]_x),\end{aligned}$$

where $4 \leq i \leq m$, is quasilinear.

- ② The equations of the extremals of \mathcal{L}_{23}^1 are given by $R^1 = \varphi^{-1}(M \times \{0\})$.
- ③ φ is a submersion. Hence, R^1 is a fibred submanifold of $J^1(FM)$.

Symbol of the PDEs system of extremals of \mathcal{L}_{23}^1

$T = TM$, $T^* = T^*M$. FM open subbundle of $\oplus^m T$.

Symbol of the PDEs system of extremals of \mathcal{L}_{23}^1

$T = TM$, $T^* = T^*M$. FM open subbundle of $\oplus^m T$.

$$\sigma: FM \times_M (T^* \otimes (\oplus^m T)) \longrightarrow M \times \mathbb{R}^{3(m-2)},$$

Symbol of the PDEs system of extremals of \mathcal{L}_{23}^1

$T = TM$, $T^* = T^*M$. FM open subbundle of $\oplus^m T$.

$$\sigma: FM \times_M (T^* \otimes (\oplus^m T)) \longrightarrow M \times \mathbb{R}^{3(m-2)},$$

$$\sigma^{i-3}(u, w \otimes v) = (w \otimes v_i)((X_3)_x, (\omega^1)_x) - (w \otimes v_3)((X_i)_x, (\omega^1)_x),$$

$$\sigma^{m-6+i}(u, w \otimes v) = (w \otimes v_i)((X_2)_x, (\omega^1)_x) - (w \otimes v_2)((X_i)_x, (\omega^1)_x),$$

$$\begin{aligned} \sigma^{2m-5}(u, w \otimes v) = & 2[(w \otimes v_3)((X_1)_x, (\omega^1)_x) - (w \otimes v_1)((X_3)_x, (\omega^1)_x)] \\ & + \sum_{b=4}^m [(w \otimes v_3)((X_b)_x, (\omega^b)_x) - (w \otimes v_b)((X_3)_x, (\omega^b)_x)], \end{aligned}$$

$$\begin{aligned} \sigma^{2m-4}(u, w \otimes v) = & 2[(w \otimes v_2)((X_1)_x, (\omega^1)_x) - (w \otimes v_1)((X_2)_x, (\omega^1)_x)] \\ & + \sum_{b=4}^m [(w \otimes v_2)((X_b)_x, (\omega^b)_x) - (w \otimes v_b)((X_2)_x, (\omega^b)_x)], \end{aligned}$$

$$\sigma^{2m-3}(u, w \otimes v) = (w \otimes v_3)((X_2)_x, (\omega^1)_x) - (w \otimes v_2)((X_3)_x, (\omega^1)_x),$$

$$\sigma^{2m-6+i}(u, w \otimes v) = (w \otimes v_3)((X_2)_x, (\omega^i)_x) - (w \otimes v_2)((X_3)_x, (\omega^i)_x),$$

Symbol of the PDEs system of extremals of \mathcal{L}_{23}^1

$T = TM$, $T^* = T^*M$. FM open subbundle of $\oplus^m T$.

$$\sigma: FM \times_M (T^* \otimes (\oplus^m T)) \longrightarrow M \times \mathbb{R}^{3(m-2)},$$

$$\sigma^{i-3}(u, w \otimes v) = (w \otimes v_i)((X_3)_x, (\omega^1)_x) - (w \otimes v_3)((X_i)_x, (\omega^1)_x),$$

$$\sigma^{m-6+i}(u, w \otimes v) = (w \otimes v_i)((X_2)_x, (\omega^1)_x) - (w \otimes v_2)((X_i)_x, (\omega^1)_x),$$

$$\begin{aligned} \sigma^{2m-5}(u, w \otimes v) = & 2[(w \otimes v_3)((X_1)_x, (\omega^1)_x) - (w \otimes v_1)((X_3)_x, (\omega^1)_x)] \\ & + \sum_{b=4}^m [(w \otimes v_3)((X_b)_x, (\omega^b)_x) - (w \otimes v_b)((X_3)_x, (\omega^b)_x)], \end{aligned}$$

$$\begin{aligned} \sigma^{2m-4}(u, w \otimes v) = & 2[(w \otimes v_2)((X_1)_x, (\omega^1)_x) - (w \otimes v_1)((X_2)_x, (\omega^1)_x)] \\ & + \sum_{b=4}^m [(w \otimes v_2)((X_b)_x, (\omega^b)_x) - (w \otimes v_b)((X_2)_x, (\omega^b)_x)], \end{aligned}$$

$$\sigma^{2m-3}(u, w \otimes v) = (w \otimes v_3)((X_2)_x, (\omega^1)_x) - (w \otimes v_2)((X_3)_x, (\omega^1)_x),$$

$$\sigma^{2m-6+i}(u, w \otimes v) = (w \otimes v_3)((X_2)_x, (\omega^i)_x) - (w \otimes v_2)((X_3)_x, (\omega^i)_x),$$

where $4 \leq i \leq m$, $w \in T_x^*$ and $v \in \oplus^m T_x$.

Cauchy-Kowaleska PDEs system

First-order PDEs system of the **Cauchy-Kowaleska type**:

$$\frac{\partial u^\beta}{\partial x^m} = \Phi^\beta \left(x^i, u^\alpha, \frac{\partial u^\alpha}{\partial x^h} \right), \quad \begin{array}{l} 1 \leq i \leq m, \\ 1 \leq h \leq m-1, \\ \alpha, \beta = 1 \dots n. \end{array}$$

Cauchy-Kowaleska PDEs system

First-order PDEs system of the **Cauchy-Kowaleska type**:

$$\frac{\partial u^\beta}{\partial x^m} = \Phi^\beta \left(x^i, u^\alpha, \frac{\partial u^\alpha}{\partial x^h} \right), \quad \begin{array}{l} 1 \leq i \leq m, \\ 1 \leq h \leq m-1, \\ \alpha, \beta = 1 \dots n. \end{array}$$

If $\Phi^\beta \in C^\omega$ on a neighbourhood of an analytic initial data then the Cauchy problem has a unique analytic solution.

Cauchy-Kowaleska PDEs system

First-order PDEs system of the **Cauchy-Kowaleska type**:

$$\frac{\partial u^\beta}{\partial x^m} = \Phi^\beta \left(x^i, u^\alpha, \frac{\partial u^\alpha}{\partial x^h} \right), \quad \begin{array}{l} 1 \leq i \leq m, \\ 1 \leq h \leq m-1, \\ \alpha, \beta = 1 \dots n. \end{array}$$

If $\Phi^\beta \in C^\omega$ on a neighbourhood of an analytic initial data then the Cauchy problem has a unique analytic solution.

$p: P \rightarrow M$ fibred manifold, $\dim P = m + n$.

Cauchy-Kowaleska PDEs system

First-order PDEs system of the **Cauchy-Kowaleska type**:

$$\frac{\partial u^\beta}{\partial x^m} = \Phi^\beta \left(x^i, u^\alpha, \frac{\partial u^\alpha}{\partial x^h} \right), \quad \begin{array}{l} 1 \leq i \leq m, \\ 1 \leq h \leq m-1, \\ \alpha, \beta = 1 \dots n. \end{array}$$

If $\Phi^\beta \in C^\omega$ on a neighbourhood of an analytic initial data then the Cauchy problem has a unique analytic solution.

$p: P \rightarrow M$ fibred manifold, $\dim P = m + n$.

Let $R^1 \subset J^1 P$ be a fibred submanifold given by

$$F^\beta(x^i, y^\alpha, y_i^\alpha) = 0, \quad 1 \leq i \leq m, \quad \alpha, \beta = 1 \dots n.$$

Cauchy-Kowaleska PDEs system

First-order PDEs system of the **Cauchy-Kowaleska type**:

$$\frac{\partial u^\beta}{\partial x^m} = \Phi^\beta \left(x^i, u^\alpha, \frac{\partial u^\alpha}{\partial x^h} \right), \quad \begin{array}{l} 1 \leq i \leq m, \\ 1 \leq h \leq m-1, \\ \alpha, \beta = 1 \dots n. \end{array}$$

If $\Phi^\beta \in C^\omega$ on a neighbourhood of an analytic initial data then the Cauchy problem has a unique analytic solution.

$p: P \rightarrow M$ fibred manifold, $\dim P = m + n$.

Let $R^1 \subset J^1 P$ be a fibred submanifold given by

$$F^\beta(x^i, y^\alpha, y_i^\alpha) = 0, \quad 1 \leq i \leq m, \quad \alpha, \beta = 1 \dots n.$$

$$M' = \{x \in M \mid x^m = 0\}.$$

Cauchy-Kowaleska PDEs system

First-order PDEs system of the **Cauchy-Kowaleska type**:

$$\frac{\partial u^\beta}{\partial x^m} = \Phi^\beta \left(x^i, u^\alpha, \frac{\partial u^\alpha}{\partial x^h} \right), \quad \begin{array}{l} 1 \leq i \leq m, \\ 1 \leq h \leq m-1, \\ \alpha, \beta = 1 \dots n. \end{array}$$

If $\Phi^\beta \in C^\omega$ on a neighbourhood of an analytic initial data then the Cauchy problem has a unique analytic solution.

$p: P \rightarrow M$ fibred manifold, $\dim P = m + n$.

Let $R^1 \subset J^1 P$ be a fibred submanifold given by

$$F^\beta(x^i, y^\alpha, y_i^\alpha) = 0, \quad 1 \leq i \leq m, \quad \alpha, \beta = 1 \dots n.$$

$M' = \{x \in M \mid x^m = 0\}$. A $x_0 \in M'$ is **non-characteristic** of R^1 if and only if the system can locally be written as a Cauchy-Kowaleska system;

Cauchy-Kowaleska PDEs system

First-order PDEs system of the **Cauchy-Kowaleska type**:

$$\frac{\partial u^\beta}{\partial x^m} = \Phi^\beta \left(x^i, u^\alpha, \frac{\partial u^\alpha}{\partial x^h} \right), \quad \begin{array}{l} 1 \leq i \leq m, \\ 1 \leq h \leq m-1, \\ \alpha, \beta = 1 \dots n. \end{array}$$

If $\Phi^\beta \in C^\omega$ on a neighbourhood of an analytic initial data then the Cauchy problem has a unique analytic solution.

$p: P \rightarrow M$ fibred manifold, $\dim P = m + n$.

Let $R^1 \subset J^1 P$ be a fibred submanifold given by

$$F^\beta(x^i, y^\alpha, y_i^\alpha) = 0, \quad 1 \leq i \leq m, \quad \alpha, \beta = 1 \dots n.$$

$M' = \{x \in M \mid x^m = 0\}$. A $x_0 \in M'$ is **non-characteristic** of R^1 if and only if the system can locally be written as a Cauchy-Kowaleska system; i.e.,

$$\det(\partial F^\beta / \partial y_m^\alpha)(j_{x_0}^1 s) \neq 0.$$

Cauchy-Kowaleska PDEs system

First-order PDEs system of the **Cauchy-Kowaleska type**:

$$\frac{\partial u^\beta}{\partial x^m} = \Phi^\beta \left(x^i, u^\alpha, \frac{\partial u^\alpha}{\partial x^h} \right), \quad \begin{array}{l} 1 \leq i \leq m, \\ 1 \leq h \leq m-1, \\ \alpha, \beta = 1 \dots n. \end{array}$$

If $\Phi^\beta \in C^\omega$ on a neighbourhood of an analytic initial data then the Cauchy problem has a unique analytic solution.

$p: P \rightarrow M$ fibred manifold, $\dim P = m + n$.

Let $R^1 \subset J^1 P$ be a fibred submanifold given by

$$F^\beta(x^i, y^\alpha, y_i^\alpha) = 0, \quad 1 \leq i \leq m, \quad \alpha, \beta = 1 \dots n.$$

$M' = \{x \in M \mid x^m = 0\}$. A $x_0 \in M'$ is **non-characteristic** of R^1 if and only if the system can locally be written as a Cauchy-Kowaleska system; i.e.,

$$\det(\partial F^\beta / \partial y_m^\alpha)(j_{x_0}^1 s) \neq 0.$$

Otherwise, x_0 is **characteristic**.

GOAL

- We extend the notion of a characteristic submanifold to the case of underdetermined PDEs systems, that is, $R^1 \subset J^1 P$ being a PDEs system of codimension $\nu \leq n$.

- We extend the notion of a characteristic submanifold to the case of underdetermined PDEs systems, that is, $R^1 \subset J^1 P$ being a PDEs system of codimension $\nu \leq n$.
- We show that every hypersurface is characteristic for the PDEs of the extremals of \mathcal{L}_{23}^1 .

- We extend the notion of a characteristic submanifold to the case of underdetermined PDEs systems, that is, $R^1 \subset J^1 P$ being a PDEs system of codimension $\nu \leq n$.
- We show that every hypersurface is characteristic for the PDEs of the extremals of \mathcal{L}_{23}^1 .
- This system is proved to be formally integrable.

Characteristic submanifold for an arbitrary PDEs system

Definition

Characteristic submanifold for an arbitrary PDEs system

Definition

$p: P \rightarrow M$ and $R^1 \subset J^1 P$ be a PDEs system of codimension $\nu \leq n$.

Characteristic submanifold for an arbitrary PDEs system

Definition

$p: P \rightarrow M$ and $R^1 \subset J^1 P$ be a PDEs system of codimension $\nu \leq n$.
Let M' be a hypersurface of M and $x_0 \in M'$.

Characteristic submanifold for an arbitrary PDEs system

Definition

$p: P \rightarrow M$ and $R^1 \subset J^1 P$ be a PDEs system of codimension $\nu \leq n$.

Let M' be a hypersurface of M and $x_0 \in M'$.

We consider

$$T_{x_0}^0 M' = \{w \in T_{x_0}^* M \mid w(\nu) = 0, \forall \nu \in T_{x_0} M'\}.$$

Characteristic submanifold for an arbitrary PDEs system

Definition

$p: P \rightarrow M$ and $R^1 \subset J^1 P$ be a PDEs system of codimension $\nu \leq n$.

Let M' be a hypersurface of M and $x_0 \in M'$.

We consider

$$T_{x_0}^0 M' = \{w \in T_{x_0}^* M \mid w(\nu) = 0, \forall \nu \in T_{x_0} M'\}.$$

A point $x_0 \in M'$ is said to be **non-characteristic** of R^1 if there exists $j_{x_0}^1 s \in R^1$ such that

$$\dim[\varepsilon(T_{x_0}^0 M' \otimes V_{s(x_0)}(p)) \cap T_{j_{x_0}^1 s} R^1] = n - \nu,$$

Characteristic submanifold for an arbitrary PDEs system

Definition

$p: P \rightarrow M$ and $R^1 \subset J^1 P$ be a PDEs system of codimension $\nu \leq n$.

Let M' be a hypersurface of M and $x_0 \in M'$.

We consider

$$T_{x_0}^0 M' = \{w \in T_{x_0}^* M \mid w(\nu) = 0, \forall \nu \in T_{x_0} M'\}.$$

A point $x_0 \in M'$ is said to be **non-characteristic** of R^1 if there exists $j_{x_0}^1 s \in R^1$ such that

$$\dim[\varepsilon(T_{x_0}^0 M' \otimes V_{s(x_0)}(p)) \cap T_{j_{x_0}^1 s} R^1] = n - \nu,$$

$V_{s(x_0)}(p)$ is the p -vertical tangent space,

Characteristic submanifold for an arbitrary PDEs system

Definition

$p: P \rightarrow M$ and $R^1 \subset J^1 P$ be a PDEs system of codimension $\nu \leq n$.
Let M' be a hypersurface of M and $x_0 \in M'$.

We consider

$$T_{x_0}^0 M' = \{w \in T_{x_0}^* M \mid w(\nu) = 0, \forall \nu \in T_{x_0} M'\}.$$

A point $x_0 \in M'$ is said to be **non-characteristic** of R^1 if there exists $j_{x_0}^1 s \in R^1$ such that

$$\dim[\varepsilon(T_{x_0}^0 M' \otimes V_{s(x_0)}(p)) \cap T_{j_{x_0}^1 s} R^1] = n - \nu,$$

$V_{s(x_0)}(p)$ is the p -vertical tangent space, and $T_{x_0}^0 M' \otimes V_{s(x_0)}(p)$ is identified with a subspace of $V_{j_{x_0}^1 s}(p_0^1)$.

Characteristic submanifold for an arbitrary PDEs system

Definition

$p: P \rightarrow M$ and $R^1 \subset J^1 P$ be a PDEs system of codimension $\nu \leq n$. Let M' be a hypersurface of M and $x_0 \in M'$.

We consider

$$T_{x_0}^0 M' = \{w \in T_{x_0}^* M \mid w(\nu) = 0, \forall \nu \in T_{x_0} M'\}.$$

A point $x_0 \in M'$ is said to be **non-characteristic** of R^1 if there exists $j_{x_0}^1 s \in R^1$ such that

$$\dim[\varepsilon(T_{x_0}^0 M' \otimes V_{s(x_0)}(p)) \cap T_{j_{x_0}^1 s} R^1] = n - \nu,$$

$V_{s(x_0)}(p)$ is the p -vertical tangent space, and $T_{x_0}^0 M' \otimes V_{s(x_0)}(p)$ is identified with a subspace of $V_{j_{x_0}^1 s}(p_0^1)$.

Recall

$$0 \rightarrow V_{j_{x_0}^1 s}(p_0^1) \xrightarrow{\varepsilon} T_{j_{x_0}^1 s}(J^1 P) \xrightarrow{(p_0^1)^*} T_{s(x_0)}(P) \rightarrow 0.$$

Characteristic submanifold for an arbitrary PDEs system

Local condition

Characteristic submanifold for an arbitrary PDEs system

Local condition

Let (x^i, y^α) be a fibred coordinate system for p such that in a neighbourhood of $x_0 \in M'$ the hypersurface M' is defined by $x^m = 0$.

Characteristic submanifold for an arbitrary PDEs system

Local condition

Let (x^i, y^α) be a fibred coordinate system for p such that in a neighbourhood of $x_0 \in M'$ the hypersurface M' is defined by $x^m = 0$. In this case, $T_{x_0}^0 M' \otimes V_{s(x_0)}(p) = \langle dx^m \otimes \partial/\partial y^\alpha \equiv \partial/\partial y_m^\alpha \rangle$.

Characteristic submanifold for an arbitrary PDEs system

Local condition

Let (x^i, y^α) be a fibred coordinate system for p such that in a neighbourhood of $x_0 \in M'$ the hypersurface M' is defined by $x^m = 0$. In this case, $T_{x_0}^0 M' \otimes V_{s(x_0)}(p) = \langle dx^m \otimes \partial/\partial y^\alpha \equiv \partial/\partial y_m^\alpha \rangle$.

The point $x_0 \in M'$ is non-characteristic of the PDEs system R^1 given by

$$F^k(x^i, y^\alpha, y_i^\alpha) = 0, \quad 1 \leq k \leq \nu,$$

Characteristic submanifold for an arbitrary PDEs system

Local condition

Let (x^i, y^α) be a fibred coordinate system for p such that in a neighbourhood of $x_0 \in M'$ the hypersurface M' is defined by $x^m = 0$.

In this case, $T_{x_0}^0 M' \otimes V_{s(x_0)}(p) = \langle dx^m \otimes \partial/\partial y^\alpha \equiv \partial/\partial y_m^\alpha \rangle$.

The point $x_0 \in M'$ is non-characteristic of the PDEs system R^1 given by

$$F^k(x^i, y^\alpha, y_i^\alpha) = 0, \quad 1 \leq k \leq \nu,$$

if and only if

$$\operatorname{rk} \left(\left(\frac{\partial F^k}{\partial y_m^\alpha} (j_{x_0}^1 s) \right)_\alpha^k \right) = \nu.$$

Characteristic submanifold for an arbitrary PDEs system

Local condition

Let (x^i, y^α) be a fibred coordinate system for p such that in a neighbourhood of $x_0 \in M'$ the hypersurface M' is defined by $x^m = 0$.

In this case, $T_{x_0}^0 M' \otimes V_{s(x_0)}(p) = \langle dx^m \otimes \partial/\partial y^\alpha \equiv \partial/\partial y_m^\alpha \rangle$.

The point $x_0 \in M'$ is non-characteristic of the PDEs system R^1 given by

$$F^k(x^i, y^\alpha, y_i^\alpha) = 0, \quad 1 \leq k \leq \nu,$$

if and only if

$$\operatorname{rk} \left(\left(\frac{\partial F^k}{\partial y_m^\alpha} (j_{x_0}^1 s) \right)_\alpha^k \right) = \nu.$$

In that case, R^1 is a **generalized Cauchy-Kowaleska** system.

Characteristics for the PDEs of extremals of \mathcal{L}_{23}^1

Characteristics for the PDEs of extremals of \mathcal{L}_{23}^1

For $R^1 = \varphi^{-1}(M \times \{0\})$, $M' = \{x_0 \in M \mid x^m = 0\}$,

Characteristics for the PDEs of extremals of \mathcal{L}_{23}^1

For $R^1 = \varphi^{-1}(M \times \{0\})$, $M' = \{x_0 \in M \mid x^m = 0\}$, we have:

$$\dim[(T_{x_0}^0 M' \otimes V_{s(x_0)}(\pi)) \cap T_{j_{x_0}^1 s} R^1] = m^2 - r,$$

Characteristics for the PDEs of extremals of \mathcal{L}_{23}^1

For $R^1 = \varphi^{-1}(M \times \{0\})$, $M' = \{x_0 \in M \mid x^m = 0\}$, we have:

$$\dim[(T_{x_0}^0 M' \otimes V_{s(x_0)}(\pi)) \cap T_{j_{x_0}^1 s} R^1] = m^2 - r,$$

where

$$r = \text{rk} \left(\left(\frac{\partial \varphi^k}{\partial x_{m,b}^a} (j_{x_0}^1 s) \right)_{a,b=1 \dots m}^{k=1 \dots 3(m-2)} \right) < 3(m-2).$$

Characteristics for the PDEs of extremals of \mathcal{L}_{23}^1

For $R^1 = \varphi^{-1}(M \times \{0\})$, $M' = \{x_0 \in M \mid x^m = 0\}$, we have:

$$\dim[(T_{x_0}^0 M' \otimes V_{s(x_0)}(\pi)) \cap T_{j_{x_0}^1 s} R^1] = m^2 - r,$$

where

$$r = \text{rk} \left(\left(\frac{\partial \varphi^k}{\partial x_{m,b}^a} (j_{x_0}^1 s) \right)_{a,b=1 \dots m}^{k=1 \dots 3(m-2)} \right) < 3(m-2).$$

Theorem

Characteristics for the PDEs of extremals of \mathcal{L}_{23}^1

For $R^1 = \varphi^{-1}(M \times \{0\})$, $M' = \{x_0 \in M \mid x^m = 0\}$, we have:

$$\dim[(T_{x_0}^0 M' \otimes V_{s(x_0)}(\pi)) \cap T_{j_{x_0}^1 s} R^1] = m^2 - r,$$

where

$$r = \text{rk} \left(\left(\frac{\partial \varphi^k}{\partial x_{m,b}^a} (j_{x_0}^1 s) \right)_{a,b=1\dots m}^{k=1\dots 3(m-2)} \right) < 3(m-2).$$

Theorem

Every hypersurface of M is characteristic for the PDEs system of the extremals of \mathcal{L}_{23}^1 .

Characteristics for the PDEs of extremals of \mathcal{L}_{23}^1

For $R^1 = \varphi^{-1}(M \times \{0\})$, $M' = \{x_0 \in M \mid x^m = 0\}$, we have:

$$\dim[(T_{x_0}^0 M' \otimes V_{s(x_0)}(\pi)) \cap T_{j_{x_0}^1 s} R^1] = m^2 - r,$$

where

$$r = \text{rk} \left(\left(\frac{\partial \varphi^k}{\partial x_{m,b}^a} (j_{x_0}^1 s) \right)_{a,b=1\dots m}^{k=1\dots 3(m-2)} \right) < 3(m-2).$$

Theorem

Every hypersurface of M is characteristic for the PDEs system of the extremals of \mathcal{L}_{23}^1 .

Hence the Euler-Lagrange equations for \mathcal{L}_{23}^1 cannot be written in the Cauchy-Kowaleska form, but they are formally integrable and consequently the Cauchy problem admits a solution.

Formal integrability (recall)

Formal integrability (recall)



R. Bryant, S. S. Chern, R. Gardner, H. Goldschmidt, and P. Griffiths, *Exterior differential systems*, Math. Sci. Res. Inst. Publ., Volume 18, Springer-Verlag, Berlin and New York, 1991.

Formal integrability (recall)



R. Bryant, S. S. Chern, R. Gardner, H. Goldschmidt, and P. Griffiths, *Exterior differential systems*, Math. Sci. Res. Inst. Publ., Volume 18, Springer-Verlag, Berlin and New York, 1991.

$p: P \rightarrow M$ fibred manifold. A PDEs system $R^k \subset J^k(P)$ is **formally integrable** if

$$p_{k+l-1}^{k+l}: R^{k+l} \longrightarrow R^{k+l-1}$$

are surjective $\forall l \geq 0$, where $R^{k+l} \subset J^{k+l}(P)$ is the l -prolongation of R^k .

Formal integrability (recall)



R. Bryant, S. S. Chern, R. Gardner, H. Goldschmidt, and P. Griffiths, *Exterior differential systems*, Math. Sci. Res. Inst. Publ., Volume 18, Springer-Verlag, Berlin and New York, 1991.

$p: P \rightarrow M$ fibred manifold. A PDEs system $R^k \subset J^k(P)$ is **formally integrable** if

$$p_{k+l-1}^{k+l}: R^{k+l} \longrightarrow R^{k+l-1}$$

are surjective $\forall l \geq 0$, where $R^{k+l} \subset J^{k+l}(P)$ is the l -prolongation of R^k .

Theorem (Goldschmidt)

If the following conditions are satisfied:

Formal integrability (recall)



R. Bryant, S. S. Chern, R. Gardner, H. Goldschmidt, and P. Griffiths, *Exterior differential systems*, Math. Sci. Res. Inst. Publ., Volume 18, Springer-Verlag, Berlin and New York, 1991.

$p: P \rightarrow M$ fibred manifold. A PDEs system $R^k \subset J^k(P)$ is **formally integrable** if

$$p_{k+l-1}^{k+l}: R^{k+l} \longrightarrow R^{k+l-1}$$

are surjective $\forall l \geq 0$, where $R^{k+l} \subset J^{k+l}(P)$ is the l -prolongation of R^k .

Theorem (Goldschmidt)

If the following conditions are satisfied:

- 1 $\mathfrak{g}_{k+1} = \ker \sigma_1$ is a vector bundle over R^k ,

Formal integrability (recall)



R. Bryant, S. S. Chern, R. Gardner, H. Goldschmidt, and P. Griffiths, *Exterior differential systems*, Math. Sci. Res. Inst. Publ., Volume 18, Springer-Verlag, Berlin and New York, 1991.

$p: P \rightarrow M$ fibred manifold. A PDEs system $R^k \subset J^k(P)$ is **formally integrable** if

$$p_{k+l-1}^{k+l}: R^{k+l} \longrightarrow R^{k+l-1}$$

are surjective $\forall l \geq 0$, where $R^{k+l} \subset J^{k+l}(P)$ is the l -prolongation of R^k .

Theorem (Goldschmidt)

If the following conditions are satisfied:

- 1 $\mathfrak{g}_{k+1} = \ker \sigma_1$ is a vector bundle over R^k ,
- 2 $p_{k-1}^k: R^k \longrightarrow R^{k-1}$ is surjective,

Formal integrability (recall)



R. Bryant, S. S. Chern, R. Gardner, H. Goldschmidt, and P. Griffiths, *Exterior differential systems*, Math. Sci. Res. Inst. Publ., Volume 18, Springer-Verlag, Berlin and New York, 1991.

$p: P \rightarrow M$ fibred manifold. A PDEs system $R^k \subset J^k(P)$ is **formally integrable** if

$$p_{k+l-1}^{k+l}: R^{k+l} \longrightarrow R^{k+l-1}$$

are surjective $\forall l \geq 0$, where $R^{k+l} \subset J^{k+l}(P)$ is the l -prolongation of R^k .

Theorem (Goldschmidt)

If the following conditions are satisfied:

- 1 $\mathfrak{g}_{k+1} = \ker \sigma_1$ is a vector bundle over R^k ,
- 2 $p_{k-1}^k: R^k \longrightarrow R^{k-1}$ is surjective,
- 3 there exists a quasi-regular basis for T_x for $\mathfrak{g}_k = \ker \sigma$ at $u \in P$,

Formal integrability (recall)



R. Bryant, S. S. Chern, R. Gardner, H. Goldschmidt, and P. Griffiths, *Exterior differential systems*, Math. Sci. Res. Inst. Publ., Volume 18, Springer-Verlag, Berlin and New York, 1991.

$p: P \rightarrow M$ fibred manifold. A PDEs system $R^k \subset J^k(P)$ is **formally integrable** if

$$p_{k+l-1}^{k+l}: R^{k+l} \longrightarrow R^{k+l-1}$$

are surjective $\forall l \geq 0$, where $R^{k+l} \subset J^{k+l}(P)$ is the l -prolongation of R^k .

Theorem (Goldschmidt)

If the following conditions are satisfied:

- ① $\mathfrak{g}_{k+1} = \ker \sigma_1$ is a vector bundle over R^k ,
- ② $p_{k-1}^k: R^k \longrightarrow R^{k-1}$ is surjective,
- ③ there exists a quasi-regular basis for T_x for $\mathfrak{g}_k = \ker \sigma$ at $u \in P$,

then, the PDEs system R^k is formally integrable.

First prolongation of the symbol

$$\sigma_1: J^1(FM) \times_M (S^2 T^* \otimes (\oplus^m T)) \longrightarrow T^* \otimes \mathbb{R}^{3(m-2)} \cong \oplus^{3(m-2)} T^*,$$

First prolongation of the symbol

$$\sigma_1: J^1(FM) \times_M (S^2 T^* \otimes (\oplus^m T)) \longrightarrow T^* \otimes \mathbb{R}^{3(m-2)} \cong \oplus^{3(m-2)} T^*,$$

$$(\sigma_1)^{i-3}(j_x^1 s, [f]_x^2 \otimes v) = (\omega^1(Y_i)d(X_3 f) - \omega^1(Y_3)d(X_i f))(x),$$

$$(\sigma_1)^{m-6+i}(j_x^1 s, [f]_x^2 \otimes v) = (\omega^1(Y_i)d(X_2 f) - \omega^1(Y_2)d(X_i f))(x),$$

$$\begin{aligned} (\sigma_1)^{2m-5}(j_x^1 s, [f]_x^2 \otimes v) &= 2[\omega^1(Y_3)d(X_1 f) - \omega^1(Y_1)d(X_3 f)](x) \\ &\quad + \sum_{b=4}^m [\omega^b(Y_3)d(X_b f) - \omega^b(Y_b)d(X_3 f)](x), \end{aligned}$$

$$\begin{aligned} (\sigma_1)^{2m-4}(j_x^1 s, [f]_x^2 \otimes v) &= 2[\omega^1(Y_2)d(X_1 f) - \omega^1(Y_1)d(X_2 f)](x) \\ &\quad + \sum_{b=4}^m [\omega^b(Y_2)d(X_b f) - \omega^b(Y_b)d(X_2 f)](x), \end{aligned}$$

$$(\sigma_1)^{2m-3}(j_x^1 s, [f]_x^2 \otimes v) = (\omega^1(Y_3)d(X_2 f) - \omega^1(Y_2)d(X_3 f))(x),$$

$$(\sigma_1)^{2m-6+i}(j_x^1 s, [f]_x^2 \otimes v) = (\omega^i(Y_3)d(X_2 f) - \omega^i(Y_2)d(X_3 f))(x),$$

First prolongation of the symbol

$$\sigma_1: J^1(FM) \times_M (S^2 T^* \otimes (\oplus^m T)) \longrightarrow T^* \otimes \mathbb{R}^{3(m-2)} \cong \oplus^{3(m-2)} T^*,$$

$$(\sigma_1)^{i-3}(j_x^1 s, [f]_x^2 \otimes v) = (\omega^1(Y_i)d(X_3 f) - \omega^1(Y_3)d(X_i f))(x),$$

$$(\sigma_1)^{m-6+i}(j_x^1 s, [f]_x^2 \otimes v) = (\omega^1(Y_i)d(X_2 f) - \omega^1(Y_2)d(X_i f))(x),$$

$$\begin{aligned} (\sigma_1)^{2m-5}(j_x^1 s, [f]_x^2 \otimes v) &= 2[\omega^1(Y_3)d(X_1 f) - \omega^1(Y_1)d(X_3 f)](x) \\ &\quad + \sum_{b=4}^m [\omega^b(Y_3)d(X_b f) - \omega^b(Y_b)d(X_3 f)](x), \end{aligned}$$

$$\begin{aligned} (\sigma_1)^{2m-4}(j_x^1 s, [f]_x^2 \otimes v) &= 2[\omega^1(Y_2)d(X_1 f) - \omega^1(Y_1)d(X_2 f)](x) \\ &\quad + \sum_{b=4}^m [\omega^b(Y_2)d(X_b f) - \omega^b(Y_b)d(X_2 f)](x), \end{aligned}$$

$$(\sigma_1)^{2m-3}(j_x^1 s, [f]_x^2 \otimes v) = (\omega^1(Y_3)d(X_2 f) - \omega^1(Y_2)d(X_3 f))(x),$$

$$(\sigma_1)^{2m-6+i}(j_x^1 s, [f]_x^2 \otimes v) = (\omega^i(Y_3)d(X_2 f) - \omega^i(Y_2)d(X_3 f))(x),$$

for $4 \leq i \leq m$, $[f]_x^2 \in S^2 T^*$ and $v = ((Y_1)_x, \dots, (Y_m)_x)$.

Theorem

Let Ω denote the curvature of $R^1 = \{j_x^1 s \in J^1(FM) \mid \varphi(j_x^1 s) = 0\}$;

Theorem

Let Ω denote the curvature of $R^1 = \{j_x^1 s \in J^1(FM) \mid \varphi(j_x^1 s) = 0\}$; i.e.,

$$\Omega: R^1 \rightarrow (T^* \otimes (M \times \mathbb{R}^{3(m-2)})) / \text{im } \sigma_1,$$

$$\begin{aligned}\Omega(j_x^1 s) &= \text{prol}_1(\varphi)(j_x^2 s) \bmod \text{im } \sigma_1 \\ &= j_x^1(\varphi \circ j^1 s) \bmod \text{im } \sigma_1,\end{aligned}$$

Theorem

Let Ω denote the curvature of $R^1 = \{j_x^1 s \in J^1(FM) \mid \varphi(j_x^1 s) = 0\}$; i.e.,

$$\Omega: R^1 \rightarrow (T^* \otimes (M \times \mathbb{R}^{3(m-2)})) / \text{im } \sigma_1,$$

$$\begin{aligned}\Omega(j_x^1 s) &= \text{prol}_1(\varphi)(j_x^2 s) \bmod \text{im } \sigma_1 \\ &= j_x^1(\varphi \circ j^1 s) \bmod \text{im } \sigma_1,\end{aligned}$$

$\text{prol}_1(\varphi): J^2(FM) \rightarrow J^1(M \times \mathbb{R}^{3(m-2)})$ being the first prolongation of φ .

Theorem

Let Ω denote the curvature of $R^1 = \{j_x^1 s \in J^1(FM) \mid \varphi(j_x^1 s) = 0\}$; i.e.,

$$\Omega: R^1 \rightarrow (T^* \otimes (M \times \mathbb{R}^{3(m-2)})) / \text{im } \sigma_1,$$

$$\begin{aligned}\Omega(j_x^1 s) &= \text{prol}_1(\varphi)(j_x^2 s) \bmod \text{im } \sigma_1 \\ &= j_x^1(\varphi \circ j^1 s) \bmod \text{im } \sigma_1,\end{aligned}$$

$\text{prol}_1(\varphi): J^2(FM) \rightarrow J^1(M \times \mathbb{R}^{3(m-2)})$ being the first prolongation of φ .

The curvature of the system $R^1 = \varphi^{-1}(M \times \{0\})$ vanishes.

Theorem

Let Ω denote the curvature of $R^1 = \{j_x^1 s \in J^1(FM) \mid \varphi(j_x^1 s) = 0\}$; i.e.,

$$\begin{aligned}\Omega: R^1 &\rightarrow (T^* \otimes (M \times \mathbb{R}^{3(m-2)})) / \text{im } \sigma_1, \\ \Omega(j_x^1 s) &= \text{prol}_1(\varphi)(j_x^2 s) \bmod \text{im } \sigma_1 \\ &= j_x^1(\varphi \circ j^1 s) \bmod \text{im } \sigma_1,\end{aligned}$$

$\text{prol}_1(\varphi): J^2(FM) \rightarrow J^1(M \times \mathbb{R}^{3(m-2)})$ being the first prolongation of φ .

The curvature of the system $R^1 = \varphi^{-1}(M \times \{0\})$ vanishes.

As $p_1^2(R^2) = \{j_x^1 s \in R^1 \mid \Omega(j_x^2 s) = 0\}$, then $p_1^2: R^2 \rightarrow R^1$ is surjective if and only if the curvature vanishes.

Quasi-regular basis

Quasi-regular basis

Let $\sigma: FM \times_M (T^* \otimes (\bigoplus^m T_x)) \longrightarrow M \times \mathbb{R}^{3(m-2)}$ be the symbol of $\varphi: J^1(FM) \rightarrow M \times \mathbb{R}^{3(m-2)}$.

Quasi-regular basis

Let $\sigma: FM \times_M (T^* \otimes (\bigoplus^m T_x)) \longrightarrow M \times \mathbb{R}^{3(m-2)}$ be the symbol of $\varphi: J^1(FM) \rightarrow M \times \mathbb{R}^{3(m-2)}$.

We set $\mathfrak{g}_1 = \ker \sigma$, $\mathfrak{g}_2 = \ker \sigma_1$. $\mathfrak{g}_{k,u}$ fibre of \mathfrak{g}_k over $u \in FM$.

Quasi-regular basis

Let $\sigma: FM \times_M (T^* \otimes (\bigoplus^m T_x)) \longrightarrow M \times \mathbb{R}^{3(m-2)}$ be the symbol of $\varphi: J^1(FM) \rightarrow M \times \mathbb{R}^{3(m-2)}$.

We set $\mathfrak{g}_1 = \ker \sigma$, $\mathfrak{g}_2 = \ker \sigma_1$. $\mathfrak{g}_{k,u}$ fibre of \mathfrak{g}_k over $u \in FM$.
 (t_1, \dots, t_m) basis of T_x , with dual basis $(\alpha^1, \dots, \alpha^m)$.

Quasi-regular basis

Let $\sigma: FM \times_M (T^* \otimes (\bigoplus^m T_x)) \longrightarrow M \times \mathbb{R}^{3(m-2)}$ be the symbol of $\varphi: J^1(FM) \rightarrow M \times \mathbb{R}^{3(m-2)}$.

We set $\mathfrak{g}_1 = \ker \sigma$, $\mathfrak{g}_2 = \ker \sigma_1$. $\mathfrak{g}_{k,u}$ fibre of \mathfrak{g}_k over $u \in FM$.
 (t_1, \dots, t_m) basis of T_x , with dual basis $(\alpha^1, \dots, \alpha^m)$.

$$T_{x, (t_1, \dots, t_j)}^* = \langle \alpha^{j+1}, \dots, \alpha^m \rangle.$$

Quasi-regular basis

Let $\sigma: FM \times_M (T^* \otimes (\oplus^m T_x)) \longrightarrow M \times \mathbb{R}^{3(m-2)}$ be the symbol of $\varphi: J^1(FM) \rightarrow M \times \mathbb{R}^{3(m-2)}$.

We set $\mathfrak{g}_1 = \ker \sigma$, $\mathfrak{g}_2 = \ker \sigma_1$. $\mathfrak{g}_{k,u}$ fibre of \mathfrak{g}_k over $u \in FM$.
 (t_1, \dots, t_m) basis of T_x , with dual basis $(\alpha^1, \dots, \alpha^m)$.

$$T_{x, (t_1, \dots, t_j)}^* = \langle \alpha^{j+1}, \dots, \alpha^m \rangle.$$

The basis (t_1, \dots, t_m) of T_x is **quasi-regular** for \mathfrak{g}_1 at $u = s(x)$ if

$$\dim \mathfrak{g}_{2,u} = \dim \mathfrak{g}_{1,u} + \sum_{j=1}^{m-1} \dim(\mathfrak{g}_{1,u} \cap (T_{x, (t_1, \dots, t_j)}^* \otimes (\oplus^m T_x))).$$

Quasi-regular basis

Let $\sigma: FM \times_M (T^* \otimes (\oplus^m T_x)) \longrightarrow M \times \mathbb{R}^{3(m-2)}$ be the symbol of $\varphi: J^1(FM) \rightarrow M \times \mathbb{R}^{3(m-2)}$.

We set $\mathfrak{g}_1 = \ker \sigma$, $\mathfrak{g}_2 = \ker \sigma_1$. $\mathfrak{g}_{k,u}$ fibre of \mathfrak{g}_k over $u \in FM$.
 (t_1, \dots, t_m) basis of T_x , with dual basis $(\alpha^1, \dots, \alpha^m)$.

$$T_{x,(t_1,\dots,t_j)}^* = \langle \alpha^{j+1}, \dots, \alpha^m \rangle.$$

The basis (t_1, \dots, t_m) of T_x is **quasi-regular** for \mathfrak{g}_1 at $u = s(x)$ if

$$\dim \mathfrak{g}_{2,u} = \dim \mathfrak{g}_{1,u} + \sum_{j=1}^{m-1} \dim(\mathfrak{g}_{1,u} \cap (T_{x,(t_1,\dots,t_j)}^* \otimes (\oplus^m T_x))).$$

Theorem

For every $u \in F_x M$, there exists a quasi-regular basis of T_x for \mathfrak{g}_1 at u .

Integrability of the Euler-Lagrange equations

Integrability of the Euler-Lagrange equations

Theorem

Integrability of the Euler-Lagrange equations

Theorem

Let $\varphi: J^1(FM) \rightarrow M \times \mathbb{R}^{3(m-2)}$ be the quasi-linear morphism such that $R^1 = \varphi^{-1}(M \times \{0\})$ is the set of extremals of \mathcal{L}_{23}^1 . Then,

Integrability of the Euler-Lagrange equations

Theorem

Let $\varphi: J^1(FM) \rightarrow M \times \mathbb{R}^{3(m-2)}$ be the quasi-linear morphism such that $R^1 = \varphi^{-1}(M \times \{0\})$ is the set of extremals of \mathcal{L}_{23}^1 . Then,

- 1 $\mathfrak{g}_2 = \ker \sigma_1$ is a vector bundle over R^1 of constant rank.

Integrability of the Euler-Lagrange equations

Theorem

Let $\varphi: J^1(FM) \rightarrow M \times \mathbb{R}^{3(m-2)}$ be the quasi-linear morphism such that $R^1 = \varphi^{-1}(M \times \{0\})$ is the set of extremals of \mathcal{L}_{23}^1 . Then,

- ① $\mathfrak{g}_2 = \ker \sigma_1$ is a vector bundle over R^1 of constant rank.
- ② The curvature of the system R^1 vanishes.

Integrability of the Euler-Lagrange equations

Theorem

Let $\varphi: J^1(FM) \rightarrow M \times \mathbb{R}^{3(m-2)}$ be the quasi-linear morphism such that $R^1 = \varphi^{-1}(M \times \{0\})$ is the set of extremals of \mathcal{L}_{23}^1 . Then,

- ① $\mathfrak{g}_2 = \ker \sigma_1$ is a vector bundle over R^1 of constant rank.
- ② The curvature of the system R^1 vanishes.
- ③ There exists a quasi-regular basis for T_x for $\mathfrak{g}_1 = \ker \sigma$ at $u \in FM$.

Integrability of the Euler-Lagrange equations

Theorem

Let $\varphi: J^1(FM) \rightarrow M \times \mathbb{R}^{3(m-2)}$ be the quasi-linear morphism such that $R^1 = \varphi^{-1}(M \times \{0\})$ is the set of extremals of \mathcal{L}_{23}^1 . Then,

- ① $\mathfrak{g}_2 = \ker \sigma_1$ is a vector bundle over R^1 of constant rank.
- ② The curvature of the system R^1 vanishes.
- ③ There exists a quasi-regular basis for T_x for $\mathfrak{g}_1 = \ker \sigma$ at $u \in FM$.

The Euler-Lagrange equations of \mathcal{L}_{23}^1 are formally integrable.

Integrability of the Euler-Lagrange equations

Theorem

Let $\varphi: J^1(FM) \rightarrow M \times \mathbb{R}^{3(m-2)}$ be the quasi-linear morphism such that $R^1 = \varphi^{-1}(M \times \{0\})$ is the set of extremals of \mathcal{L}_{23}^1 . Then,

- ① $\mathfrak{g}_2 = \ker \sigma_1$ is a vector bundle over R^1 of constant rank.
- ② The curvature of the system R^1 vanishes.
- ③ There exists a quasi-regular basis for T_x for $\mathfrak{g}_1 = \ker \sigma$ at $u \in FM$.

The Euler-Lagrange equations of \mathcal{L}_{23}^1 are formally integrable.

Corollary

Integrability of the Euler-Lagrange equations

Theorem

Let $\varphi: J^1(FM) \rightarrow M \times \mathbb{R}^{3(m-2)}$ be the quasi-linear morphism such that $R^1 = \varphi^{-1}(M \times \{0\})$ is the set of extremals of \mathcal{L}_{23}^1 . Then,

- 1 $\mathfrak{g}_2 = \ker \sigma_1$ is a vector bundle over R^1 of constant rank.
- 2 The curvature of the system R^1 vanishes.
- 3 There exists a quasi-regular basis for T_x for $\mathfrak{g}_1 = \ker \sigma$ at $u \in FM$.

The Euler-Lagrange equations of \mathcal{L}_{23}^1 are formally integrable.

Corollary

Given a point $j_{x_0}^1 s_0 \in R^1$, there exists a section $s: U \rightarrow FM$ of class C^∞ on a neighbourhood of x_0 such that:

Integrability of the Euler-Lagrange equations

Theorem

Let $\varphi: J^1(FM) \rightarrow M \times \mathbb{R}^{3(m-2)}$ be the quasi-linear morphism such that $R^1 = \varphi^{-1}(M \times \{0\})$ is the set of extremals of \mathcal{L}_{23}^1 . Then,

- 1 $\mathfrak{g}_2 = \ker \sigma_1$ is a vector bundle over R^1 of constant rank.
- 2 The curvature of the system R^1 vanishes.
- 3 There exists a quasi-regular basis for T_x for $\mathfrak{g}_1 = \ker \sigma$ at $u \in FM$.

The Euler-Lagrange equations of \mathcal{L}_{23}^1 are formally integrable.

Corollary

Given a point $j_{x_0}^1 s_0 \in R^1$, there exists a section $s: U \rightarrow FM$ of class C^∞ on a neighbourhood of x_0 such that:

- s is a solution to R^1 ; i.e., $j_x^1 s \in R^1, \forall x \in U$,

Integrability of the Euler-Lagrange equations

Theorem

Let $\varphi: J^1(FM) \rightarrow M \times \mathbb{R}^{3(m-2)}$ be the quasi-linear morphism such that $R^1 = \varphi^{-1}(M \times \{0\})$ is the set of extremals of \mathcal{L}_{23}^1 . Then,

- ① $\mathfrak{g}_2 = \ker \sigma_1$ is a vector bundle over R^1 of constant rank.
- ② The curvature of the system R^1 vanishes.
- ③ There exists a quasi-regular basis for T_x for $\mathfrak{g}_1 = \ker \sigma$ at $u \in FM$.

The Euler-Lagrange equations of \mathcal{L}_{23}^1 are formally integrable.

Corollary

Given a point $j_{x_0}^1 s_0 \in R^1$, there exists a section $s: U \rightarrow FM$ of class C^∞ on a neighbourhood of x_0 such that:

- s is a solution to R^1 ; i.e., $j_x^1 s \in R^1, \forall x \in U$,
- $j_{x_0}^1 s = j_{x_0}^1 s_0$.

Integrability of the Jacobi equations

Let $\mathcal{S} \subset \Gamma(M, FM)$ be the set of extremals of \mathcal{L}_{23}^1 .

Integrability of the Jacobi equations

Let $\mathcal{S} \subset \Gamma(M, FM)$ be the set of extremals of \mathcal{L}_{23}^1 . The Jacobi fields are the solutions to the linearized Hamilton-Cartan equation;

Integrability of the Jacobi equations

Let $\mathcal{S} \subset \Gamma(M, FM)$ be the set of extremals of \mathcal{L}_{23}^1 . The Jacobi fields are the solutions to the linearized Hamilton-Cartan equation; that is, a Jacobi field along an extremal $s \in \mathcal{S}$ is a π -vertical vector field defined along s , $X \in \Gamma(M, s^*V(\pi))$, satisfying the Jacobi equation

Integrability of the Jacobi equations

Let $\mathcal{S} \subset \Gamma(M, FM)$ be the set of extremals of \mathcal{L}_{23}^1 . The Jacobi fields are the solutions to the linearized Hamilton-Cartan equation; that is, a Jacobi field along an extremal $s \in \mathcal{S}$ is a π -vertical vector field defined along s , $X \in \Gamma(M, s^*V(\pi))$, satisfying the Jacobi equation

$$(j^1 s)^*(i_Y L_{X(1)} d\Theta) = 0, \quad \forall Y \in \mathfrak{X}(J^1(FM)).$$

Integrability of the Jacobi equations

Let $\mathcal{S} \subset \Gamma(M, FM)$ be the set of extremals of \mathcal{L}_{23}^1 . The Jacobi fields are the solutions to the linearized Hamilton-Cartan equation; that is, a Jacobi field along an extremal $s \in \mathcal{S}$ is a π -vertical vector field defined along s , $X \in \Gamma(M, s^*V(\pi))$, satisfying the Jacobi equation

$$(j^1 s)^*(i_Y L_{X(1)} d\Theta) = 0, \quad \forall Y \in \mathfrak{X}(J^1(FM)).$$

Theorem

The Jacobi equations along an arbitrary extremal $s: M \rightarrow FM$ of \mathcal{L}_{23}^1 are formally integrable.

Integrability of the Jacobi equations




Let $\mathcal{S} \subset \Gamma(M, FM)$ be the set of extremals of \mathcal{L}_{23}^1 . The Jacobi fields are the solutions to the linearized Hamilton-Cartan equation; that is, a Jacobi field along an extremal $s \in \mathcal{S}$ is a π -vertical vector field defined along s , $X \in \Gamma(M, s^*V(\pi))$, satisfying the Jacobi equation

$$(j^1 s)^*(i_Y L_{X(1)} d\Theta) = 0, \quad \forall Y \in \mathfrak{X}(J^1(FM)).$$

Theorem

The Jacobi equations along an arbitrary extremal $s: M \rightarrow FM$ of \mathcal{L}_{23}^1 are formally integrable.

Hence, if M and s are of class C^ω , then given a point $j_{x_0}^1 \tilde{\zeta}_0 \in R_s^1$ there exists a Jacobi field $X_{\tilde{\zeta}}$ defined along s on a neighbourhood of x_0 such that $j_{x_0}^1 \tilde{\zeta} = j_{x_0}^1 \tilde{\zeta}_0$.

-  R. Bryant, S. S. Chern, R. Gardner, H. Goldschmidt, and P. Griffiths, *Exterior differential systems*, Math. Sci. Res. Inst. Publ., Volume 18, Springer-Verlag, Berlin and New York, 1991.
-  J. Muñoz Masqué, M. E. Rosado, *Invariant variational problems on linear frame bundles*, J. Phys. A **35** (2002), 2013–2036.
-  J. Muñoz Masqué, M. E. Rosado, *Integrability of the field equations of invariant variational problems on linear frame bundles*, J. Geom. Phys. **49** (2004), 119–155.

Prolongations

Prolongations

There exists a quasi-linear morphism

$$\begin{aligned}\mathrm{prol}_r(\varphi): J^{1+r}(FM) &\rightarrow J^r(M \times \mathbb{R}^{3(m-2)}), \\ \mathrm{prol}_r(\varphi)(j_x^{1+r}s) &= j_x^r(\varphi \circ j^1s),\end{aligned}$$

r -th prolongation of φ ,

Prolongations

There exists a quasi-linear morphism

$$\begin{aligned}\mathrm{prol}_r(\varphi): J^{1+r}(FM) &\rightarrow J^r(M \times \mathbb{R}^{3(m-2)}), \\ \mathrm{prol}_r(\varphi)(j_x^{1+r}s) &= j_x^r(\varphi \circ j_x^1s),\end{aligned}$$

r -th prolongation of φ , whose symbol

$$\sigma(\mathrm{prol}_r(\varphi)): J^r(FM) \times_M (S^{1+r}T^* \otimes (\oplus^m T_x)) \rightarrow J^r(M \times \mathbb{R}^{3(m-2)})$$

is determined by the r -th prolongation of the symbol of φ , as

$$\sigma(\mathrm{prol}_r(\varphi))(j_x^r s, v) \in S^r T^* \otimes (M \times \mathbb{R}^{3(m-2)}), \quad \forall j_x^r s \in J^r(FM).$$

Prolongations

There exists a quasi-linear morphism

$$\begin{aligned}\mathrm{prol}_r(\varphi): J^{1+r}(FM) &\rightarrow J^r(M \times \mathbb{R}^{3(m-2)}), \\ \mathrm{prol}_r(\varphi)(j_x^{1+r}s) &= j_x^r(\varphi \circ j_x^1 s),\end{aligned}$$

r -th prolongation of φ , whose symbol

$$\sigma(\mathrm{prol}_r(\varphi)): J^r(FM) \times_M (S^{1+r}T^* \otimes (\oplus^m T_x)) \rightarrow J^r(M \times \mathbb{R}^{3(m-2)})$$

is determined by the r -th prolongation of the symbol of φ , as

$\sigma(\mathrm{prol}_r(\varphi))(j_x^r s, v) \in S^r T^* \otimes (M \times \mathbb{R}^{3(m-2)}), \forall j_x^r s \in J^r(FM)$. That is,

$$\sigma(\mathrm{prol}_r(\varphi)) = \varepsilon' \circ \sigma_r$$

Prolongations

There exists a quasi-linear morphism

$$\begin{aligned}\mathrm{prol}_r(\varphi): J^{1+r}(FM) &\rightarrow J^r(M \times \mathbb{R}^{3(m-2)}), \\ \mathrm{prol}_r(\varphi)(j_x^{1+r}s) &= j_x^r(\varphi \circ j^1s),\end{aligned}$$

r -th prolongation of φ , whose symbol

$$\sigma(\mathrm{prol}_r(\varphi)): J^r(FM) \times_M (S^{1+r}T^* \otimes (\oplus^m T_x)) \rightarrow J^r(M \times \mathbb{R}^{3(m-2)})$$

is determined by the r -th prolongation of the symbol of φ , as

$\sigma(\mathrm{prol}_r(\varphi))(j_x^r s, v) \in S^r T^* \otimes (M \times \mathbb{R}^{3(m-2)}), \forall j_x^r s \in J^r(FM)$. That is,

$$\sigma(\mathrm{prol}_r(\varphi)) = \varepsilon' \circ \sigma_r$$

$\varepsilon': S^r T^* \otimes (M \times \mathbb{R}^{3(m-2)}) \rightarrow J^r(M \times \mathbb{R}^{3(m-2)})$ being the natural injection

Prolongations

There exists a quasi-linear morphism

$$\begin{aligned}\mathrm{prol}_r(\varphi): J^{1+r}(FM) &\rightarrow J^r(M \times \mathbb{R}^{3(m-2)}), \\ \mathrm{prol}_r(\varphi)(j_x^{1+r}s) &= j_x^r(\varphi \circ j^1s),\end{aligned}$$

r -th prolongation of φ , whose symbol

$$\sigma(\mathrm{prol}_r(\varphi)): J^r(FM) \times_M (S^{1+r}T^* \otimes (\oplus^m T_x)) \rightarrow J^r(M \times \mathbb{R}^{3(m-2)})$$

is determined by the r -th prolongation of the symbol of φ , as

$\sigma(\mathrm{prol}_r(\varphi))(j_x^r s, v) \in S^r T^* \otimes (M \times \mathbb{R}^{3(m-2)}), \forall j_x^r s \in J^r(FM)$. That is,

$$\sigma(\mathrm{prol}_r(\varphi)) = \varepsilon' \circ \sigma_r$$

$\varepsilon': S^r T^* \otimes (M \times \mathbb{R}^{3(m-2)}) \rightarrow J^r(M \times \mathbb{R}^{3(m-2)})$ being the natural injection and

$$\sigma_r: J^r(FM) \times_M (S^{1+r}T^* \otimes (\oplus^m T_x)) \rightarrow S^r T^* \otimes (M \times \mathbb{R}^{3(m-2)}).$$

Quasi-regular basis

Quasi-regular basis

$p: P \longrightarrow M$, fibred manifold.

Quasi-regular basis

$p: P \longrightarrow M$, fibred manifold.

$R^k \subset J^k(P)$ PDEs system and σ symbol of R^k .

Quasi-regular basis

$p: P \longrightarrow M$, fibred manifold.

$R^k \subset J^k(P)$ PDEs system and σ symbol of R^k .

We set $\mathfrak{g}_k = \ker \sigma$, $\mathfrak{g}_{k+1} = \ker(\sigma_1)$.

Quasi-regular basis

$p: P \longrightarrow M$, fibred manifold.

$R^k \subset J^k(P)$ PDEs system and σ symbol of R^k .

We set $\mathfrak{g}_k = \ker \sigma$, $\mathfrak{g}_{k+1} = \ker(\sigma_1)$.

$\mathfrak{g}_{k,u}$ fibre of \mathfrak{g}_k over $u \in P$.

Quasi-regular basis

$p: P \longrightarrow M$, fibred manifold.

$R^k \subset J^k(P)$ PDEs system and σ symbol of R^k .

We set $\mathfrak{g}_k = \ker \sigma$, $\mathfrak{g}_{k+1} = \ker(\sigma_1)$.

$\mathfrak{g}_{k,u}$ fibre of \mathfrak{g}_k over $u \in P$.

(t_1, \dots, t_m) basis of T_x , with dual basis $(\alpha^1, \dots, \alpha^m)$.

Quasi-regular basis

$p: P \longrightarrow M$, fibred manifold.

$R^k \subset J^k(P)$ PDEs system and σ symbol of R^k .

We set $\mathfrak{g}_k = \ker \sigma$, $\mathfrak{g}_{k+1} = \ker(\sigma_1)$.

$\mathfrak{g}_{k,u}$ fibre of \mathfrak{g}_k over $u \in P$.

(t_1, \dots, t_m) basis of T_x , with dual basis $(\alpha^1, \dots, \alpha^m)$.

$$S^k T_{x, (t_1, \dots, t_j)}^* = \langle \alpha^{i_1} \odot \dots \odot \alpha^{i_k}, \quad j+1 \leq i_1 \leq \dots \leq i_k \leq m \rangle.$$

Quasi-regular basis

$p: P \longrightarrow M$, fibred manifold.

$R^k \subset J^k(P)$ PDEs system and σ symbol of R^k .

We set $\mathfrak{g}_k = \ker \sigma$, $\mathfrak{g}_{k+1} = \ker(\sigma_1)$.

$\mathfrak{g}_{k,u}$ fibre of \mathfrak{g}_k over $u \in P$.

(t_1, \dots, t_m) basis of T_x , with dual basis $(\alpha^1, \dots, \alpha^m)$.

$$S^k T_{x, (t_1, \dots, t_j)}^* = \langle \alpha^{i_1} \odot \dots \odot \alpha^{i_k}, \quad j+1 \leq i_1 \leq \dots \leq i_k \leq m \rangle.$$

The basis (t_1, \dots, t_m) of T_x is *quasi-regular* for \mathfrak{g}_k at $u = s(x)$ if

$$\dim \mathfrak{g}_{k+1,u} = \dim \mathfrak{g}_{k,u} + \sum_{j=1}^{m-1} \dim(\mathfrak{g}_{k,u} \cap (S^k T_{x, (t_1, \dots, t_j)}^* \otimes (\oplus^m T_x))).$$